

Compactly supported cohomology of buildings

M.W. Davis* J. Dymara† T. Januszkiewicz‡ J. Meier§
B. Okun

June 13, 2008

Abstract

We compute the compactly supported cohomology of the standard realization of any locally finite building.

AMS classification numbers. Primary: 20F65
Secondary: 20E42, 20F55, 20J06, 57M07.

Keywords: Building, cohomology of groups, Coxeter group.

Introduction

A building consists of a set Φ (the elements of which are called “chambers”) together with a family of equivalence relations (“adjacency relations”) on Φ indexed by a set S and a “ W -valued distance function,” $\Phi \times \Phi \rightarrow W$, where W is a Coxeter group with fundamental set of generators S . So, associated to any building there is a Coxeter system (W, S) , its *type*.

There is a construction which associates a topological space to Φ . This construction admits some freedom of choice. The idea is to choose a space X as a “model chamber” and then glue together copies of it, one for each element of Φ . To do this, it is first necessary to choose a family of closed subspaces $\{X_s\}_{s \in S}$ so that copies of X corresponding to s -adjacent chambers are glued together along X_s . (We call such a family, $\{X_s\}_{s \in S}$, a “mirror structure” on X .) Let $\mathcal{U}(\Phi, X)$ denote the topological realization of Φ where each chamber is realized by a copy of the model chamber X . (Details are given in Section 2.)

*The first author was partially supported by NSF grant DMS 0706259.

†The second author was partially supported by KBN grant N201 012 32/0718

‡The third author also was partially supported by NSF grant DMS 0706259.

§The fourth author was partially supported by a Richard King Mellon research grant.

Classically, interest has centered on buildings of spherical or affine type, meaning that W is a spherical or Euclidean reflection group, respectively. For example, each algebraic group over a local field has a corresponding affine building. However, we are mainly interested in buildings which are not classical in that their associated Coxeter systems are neither spherical nor affine. This is a large class of spaces, many of which have a great deal of symmetry. For example, such buildings arise in the theory of Kac-Moody groups (e.g., see [6, 21, 22]). Also, nonclassical buildings associated to arbitrary right-angled Coxeter groups have been a subject of recent interest in geometric group theory (e.g., see [8, 19, 26]).

Two choices for a model chamber X stand out. The first is $X = \Delta$, a simplex of dimension $\text{Card}(S) - 1$, with its codimension one faces indexed by S . This was Tits' original choice (cf. [1]). We call $\mathcal{U}(\Phi, \Delta)$ the "classical realization" of Φ . The other choice for X is the "Davis chamber" K , defined as the geometric realization of the poset \mathcal{S} of spherical subsets of S (see [9, Chapters 7, 18]). $\mathcal{U}(\Phi, K)$ is the "standard realization" of Φ . Both realizations are contractible. The standard realization is important in geometric group theory, the reason being that in this field one is interested in discrete group actions which are both proper and cocompact and these conditions are more likely to hold for the action of a group of automorphisms on the standard realization than on the classical realization. (If Φ has finite thickness, $\mathcal{U}(\Phi, K)$ is locally finite, while $\mathcal{U}(\Phi, \Delta)$ need not be.)

If a discrete group Γ acts properly and cocompactly on a locally finite, contractible CW complex Y , then the compactly supported cohomology of Y is isomorphic to the cohomology of Γ with $\mathbf{Z}\Gamma$ coefficients. In particular, it determines the virtual cohomological dimension of Γ , as well as, the number of ends of Γ , and it determines if Γ is a duality group. (For more information, see [17, Part IV].) However, as we will explain, even if one is only interested in cohomological computations in the case of the standard realization of Φ , it is necessary to carry out similar computations for various other realizations, in particular, for the classical realization.

In the classical case of an (irreducible) affine building, the two notions of model chamber agree: $\Delta = K$. So, in the affine case the study of the cohomology of cocompact lattices in $\text{Aut}(\Phi)$ is closely tied to the study of the cohomological properties of $\mathcal{U}(\Phi, \Delta)$. For example, in [2] Borel and Serre calculated the compactly supported cohomology, $H_c^*(\mathcal{U}(\Phi, \Delta))$, for any (irreducible) affine building and then used this calculation to derive information about the cohomology of " S -arithmetic" subgroups. The calculation of [2] was that $H_c^*(\mathcal{U}(\Phi, \Delta))$ is concentrated in the top degree ($= \dim \Delta$) and is free abelian in that degree.

Our main result, Corollary 8.2, is a calculation of $H_c^*(\mathcal{U}(\Phi, K))$, generalizing the theorem of Borel-Serre. In the case where $\Phi = W$, this was done in [7, 10, 12]. For a general (thick) building, in the case where (W, S) is right-angled, it was done in [10, Theorem 6.6]. It was claimed in full generality in [12]; however, there is a mistake in the proof (see [13]).

In order to write the formula, we need more notation. Let A be the free abelian group of finitely supported, \mathbf{Z} -valued functions on Φ . For each subset $T \subseteq S$, let A^T denote the subgroup of all functions $f \in A$ which are constant on each residue of type T . (A “residue” of type T is a certain kind of subset of Φ ; in the case of the building W , a residue of type T is a left coset of W_T , the subgroup of W generated by T .) N.B. The empty set, \emptyset , is a spherical subset and a residue of type \emptyset is just a single chamber; hence, $A^\emptyset = A$. If $U \supset T$, then $A^U \subset A^T$. Let $A^{>T} \subset A^T$ denote the \mathbf{Z} -submodule, $\sum_{U \supset T} A^U$. (Throughout this paper we will use the convention that \subseteq denotes containment and \subset will be reserved for strict containment. Also, we will use the symbol \sum for an internal sum of modules and \bigoplus to mean either an external direct sum or an internal sum which we have proved is direct.)

We shall show in Section 7 that $A^{>T}$ is a direct summand of A^T . Let \hat{A}^T be a complementary summand. As in [10] the main computation is a consequence the following Decomposition Theorem (proved as Theorem 7.4 in Section 7).

The Decomposition Theorem.

$$A = \bigoplus_{U \in \mathcal{S}} \hat{A}^U \quad \text{and, in fact, for any } T \in \mathcal{S}, \quad A^T = \bigoplus_{U \supseteq T} \hat{A}^U.$$

The point is that this theorem provides a decomposition of a coefficient system which can be used to calculate the compactly supported cohomology of any of the various realizations of Φ . The calculation in which we are most interested is the following (proved as Corollary 8.2 in Section 8).

The Main Theorem. *Suppose Φ is a building of finite thickness and type (W, S) . Let K be the geometric realization of the poset \mathcal{S} of spherical subsets of S . Then*

$$H_c^*(\mathcal{U}(\Phi, K)) \cong \bigoplus_{T \in \mathcal{S}} H^*(K, K^{S-T}) \otimes \hat{A}^T.$$

(For any subset U of S , K^U denotes the union of the K_s , with $s \in U$.)

The above theorem applies to all buildings. A general building Φ will not be highly symmetric, in that its automorphism group, $\text{Aut}(\Phi)$, can have

infinitely many orbits of chambers. However, if $\Phi/\text{Aut}(\Phi)$ is finite and if Γ is a torsion-free cocompact lattice in $\text{Aut}(\Phi)$, then the Main Theorem implies that the cohomological dimension of Γ is equal to the virtual cohomological dimension of the corresponding Coxeter group. Moreover, this dimension is

$$\text{cd}(\Gamma) = \text{vcd}(W) = \max\{k \mid H^k(K, K^{S-T}) \neq 0, \text{ for some } T \in \mathcal{S}\}$$

(cf. Corollary 9.4). As another example, such a torsion-free cocompact lattice is an n -dimensional duality group if and only if for each $T \in \mathcal{S}$, $H^*(K, K^{S-T})$ is free abelian and concentrated in degree n (cf. Corollary 9.5).

The central objective of [10] was to calculate $H_c^*(\mathcal{U}(W, K))$ as a W -module. In that paper we showed there is a filtration of $H_c^*(\mathcal{U}(W, K))$ by W -submodules so that the associated graded terms look like the terms on the right hand side of the formula in the Main Theorem. Similarly, one can ask about the G -module structure of $H_c^*(\mathcal{U}(\Phi, K))$ for any subgroup $G \subseteq \text{Aut}(\Phi)$. The methods of [10] are well adapted to the present paper. In particular, for each $T \in \mathcal{S}$, the free abelian group A^T is a G -module, as is its quotient $D^T := A^T/A^{>T}$. So, as in [10], there is a filtration of $H_c^*(\mathcal{U}(\Phi, K))$ by G -submodules and we get the following (proved as Theorem 9.1).

Theorem. *Suppose G is a group of automorphisms of Φ . There is a filtration of $H_c^*(\mathcal{U}(\Phi, K))$ by right G -submodules with associated graded term in filtration degree p :*

$$\bigoplus_{\substack{T \in \mathcal{S} \\ |T|=p}} H^*(K, K^{S-T}) \otimes D^T.$$

As we mentioned earlier, although the Main Theorem is the result of importance in geometric group theory, it is no harder to do similar calculations when the chambers are modeled on an arbitrary X (or at least on X^f , the complement of the faces of X which have infinite stabilizers in W). In fact, as we explain below, the proof of the Decomposition Theorem depends on first establishing a version of the Main Theorem for $\mathcal{U}(\Phi, \Delta^f)$. So, the Main Theorem ultimately depends on first proving a version of it in the case of the classical realization. This version (proved as Theorem 4.3) is the following.

Theorem. *When W infinite, $H_c^*(\mathcal{U}(\Phi, \Delta^f))$ is free abelian and is concentrated in the top degree n ($= \dim \Delta$).*

This result is obvious when $\Phi = W$, for then $\mathcal{U}(W, \Delta^f)$ is homeomorphic to Euclidean space \mathbf{R}^n (see Section 2). We prove it for a general Φ by showing the following (Theorem 3.1).

Theorem. $\mathcal{U}(\Phi, \Delta^f)$ admits a CAT(0) metric (extending Moussong’s CAT(0) metric on the standard realization).

The existence of this CAT(0) metric on $\mathcal{U}(\Phi, \Delta^f)$ is of independent interest. (See [20, 8] or [9] for a description of Moussong’s metric on $\mathcal{U}(\Phi, K)$.)

To finish the calculation of $H_c^*(\mathcal{U}(\Phi, \Delta^f))$ we invoke a result of [4] which asserts that the compactly supported cohomology of such a CAT(0) space is concentrated in the top degree provided the cohomology of each “punctured link” vanishes except in the top degree. These links are spherical buildings and the vanishing of the cohomology groups, in degrees below the top, of their punctured versions is a result of [16] (and independently, [24]).

As we have said, special cases of the Decomposition Theorem were proved in [10]. The method of [10] was simply to find a basis for A adapted to its decomposition into the \hat{A}^T . In the general case, finding an explicit description of such a basis seems problematic. We use instead an idea coming from an analogy with the argument of [11]. In that paper we proved L^2 versions of the Decomposition Theorem and of the Main Theorem. The proof of the L^2 version of the Decomposition Theorem was homological: the key step was to show that certain “weighted L^2 -homology” groups of certain auxiliary spaces associated to W vanished except in the bottom degree. We then applied a certain duality (not applicable here) to deduce the Decomposition Theorem in the cases of actual interest. Analogously, in this paper we prove the Decomposition Theorem by establishing the vanishing, except in the top degree, of the cohomology of certain auxiliary spaces. The most important of these auxiliary spaces is $\mathcal{U}(\Phi, \Delta^f)$ and we indicated in the previous paragraphs how we prove the result in that case.

1 Coxeter groups and buildings

A *chamber system* over a set S is a set Φ of *chambers* together with a family of equivalence relations on Φ indexed by S . Two chambers are *s-equivalent* if they are related via the equivalence relation with index s ; they are *s-adjacent* if they are *s-equivalent* and not equal. A *gallery* in Φ is a finite sequence of chambers $(\varphi_0, \dots, \varphi_k)$ such that φ_{j-1} is adjacent to φ_j , $1 \leq j \leq k$. The *type* of this gallery is the word $\mathbf{s} = (s_1, \dots, s_k)$ where φ_{j-1} is s_j -adjacent to φ_j . If each s_j belongs to a given subset T of S , then the gallery is a *T-gallery*. A chamber system is *connected* (resp., *T-connected*) if any two chambers can be joined by a gallery (resp., a *T-gallery*). The *T-connected* components of a chamber system Φ are its *residues* of *type T*.

A *Coxeter matrix* over a set S is an $S \times S$ symmetric matrix $M = (m_{st})$ with each diagonal entry $= 1$ and each off-diagonal entry an integer ≥ 2 or the symbol ∞ . The matrix M defines a presentation of a group W as follows: the set of generators is S and the relations have the form $(st)^{m_{st}}$ where (s, t) ranges over all pairs in $S \times S$ such that $m_{st} \neq \infty$. The pair (W, S) is a *Coxeter system* (cf. [3, 9]). Given $T \subseteq S$, W_T denotes the subgroup generated by T ; it is called a *special subgroup*. (W_T, T) is itself a Coxeter system. The subset T is *spherical* if W_T is finite.

Definition 1.1. The poset of spherical subsets of S (partially ordered by inclusion) is denoted \mathcal{S} .

Also, \mathcal{P} ($= \mathcal{P}(S)$) will denote the poset of all proper subsets of S . (In Section 3 the poset $\mathcal{P} - \mathcal{S}$ plays a role.)

Suppose (W, S) is a Coxeter system and $M = (m_{st})$ is its Coxeter matrix. Following [23] (or [10]), a *building of type (W, S)* (or of *type M*) is a chamber system Φ over S such that

- (i) for all $s \in S$, each s -equivalence class contains at least two chambers, and
- (ii) there exists a W -valued distance function $\delta : \Phi \times \Phi \rightarrow W$. (This means that given a reduced word \mathbf{s} for an element $w \in W$, chambers φ and φ' can be joined by a gallery of type \mathbf{s} from φ to φ' if and only if $\delta(\varphi, \varphi') = w$.)

Example 1.2. The group W itself has the structure of a building: the s -equivalence classes are the left cosets of $W_{\{s\}}$ and the W -valued distance, $\delta : W \times W \rightarrow W$, is defined by $\delta(v, w) = v^{-1}w$.

A residue of type T is a building; its type is (W_T, T) . A building of type (W, S) is *spherical* if W is finite. A building has *finite thickness* if each s -equivalence class is finite, for each $s \in S$. (This implies all spherical residues are finite.) *Henceforth, all buildings will be assumed to have finite thickness.*

2 Geometric realizations of Coxeter groups and buildings

A *mirror structure* over a set S on a space X is a family of subspaces $(X_s)_{s \in S}$ indexed by S . Given a mirror structure on X , a subspace $Y \subseteq X$ inherits a

mirror structure by $Y_s := Y \cap X_s$. If X is a CW complex and each X_s is a subcomplex, then X is a *mirrored CW complex*. For each nonempty subset $T \subseteq S$, define subspaces X_T and X^T by

$$X_T := \bigcap_{s \in T} X_s \quad \text{and} \quad X^T := \bigcup_{s \in T} X_s. \quad (2.1)$$

Put $X_\emptyset := X$ and $X^\emptyset := \emptyset$. Given a cell c of (a CW complex) X or a point $x \in X$, put

$$S(c) := \{s \in S \mid c \subseteq X_s\},$$

$$S(x) := \{s \in S \mid x \in X_s\}.$$

Suppose now that S is the set of generators for a Coxeter system (W, S) . Let $\Upsilon(X)$ denote the union of the nonspherical faces of X and X^f its complement in X , i.e.,

$$\Upsilon(X) := \bigcup_{T \notin \mathcal{S}} X_T \quad \text{and} \quad X^f := X - \Upsilon(X). \quad (2.2)$$

The mirror structure is *W-finite* if $\Upsilon(X) = \emptyset$.

Given a building Φ of type (W, S) and a mirrored space X over S , define an equivalence relation \sim on $\Phi \times X$ by $(\varphi, x) \sim (\varphi', x')$ if and only if $x = x'$ and $\delta(\varphi, \varphi') \in W_{S(x)}$ (i.e., φ and φ' belong to the same $S(x)$ -residue). The *X-realization* of Φ , denoted $\mathcal{U}(\Phi, X)$, is defined by

$$\mathcal{U}(\Phi, X) := (\Phi \times X) / \sim. \quad (2.3)$$

(Φ has the discrete topology.) Suppose X is a mirrored CW complex and that we are given a cell c of X and a chamber $\varphi \in \Phi$. Then $\varphi \cdot c$ denotes the corresponding cell in $\mathcal{U}(\Phi, X)$. Let $\mathcal{U}^{(i)}$ denote the set of i -cells in $\mathcal{U}(\Phi, X)$. Each such cell has the form $\varphi \cdot c$ for some $\varphi \in \Phi$ and i -cell c of X .

The classical realization. Δ denotes the simplex of dimension $|S| - 1$, with its codimension one faces indexed by S . In other words, the mirror Δ_s is a codimension one face and Δ_T (the intersection of the Δ_s over all $s \in T$) is a face of codimension $|T|$.

The simplicial complex $\mathcal{U}(W, \Delta)$ is the *Coxeter complex* of (W, S) while $\mathcal{U}(\Phi, \Delta)$ is the *classical realization* of the building Φ .

Tits constructed a representation of W on \mathbf{R}^S called “the contragredient of the canonical representation” in [3] and the “geometric representation” in [9]. The elements of S are represented by reflections across the codimension one faces of a simplicial cone C . The union of translates of C is

denoted by WC . It is a convex cone and W acts properly on its interior \mathcal{I} . If W is infinite, WC is a proper cone. If C^f denotes the complement of the nonspherical faces of C , then \mathcal{I} is equivariantly homeomorphic to $\mathcal{U}(W, C^f)$. Assume W is infinite. Then the image of $C - 0$ in projective space can be identified with the simplex Δ obtained by intersecting C with some affine hyperplane; moreover, Δ^f is identified with the intersection of C^f and this hyperplane. The image of \mathcal{I} in projective space is then identified with $\mathcal{U}(W, \Delta^f)$. Since this image is the interior of a topological disk (of dimension $|S| - 1$), it follows that $\mathcal{U}(W, \Delta^f)$ is homeomorphic to a Euclidean space of that dimension. In particular, it is contractible.

Geometric realizations of posets. Given a poset \mathcal{T} , $\text{Flag}(\mathcal{T})$ denotes the set of finite chains in \mathcal{T} , partially ordered by inclusion, i.e., an element of $\text{Flag}(\mathcal{T})$ is a finite, nonempty, totally ordered subset of \mathcal{T} . If $\alpha = \{t_0, \dots, t_k\} \in \text{Flag}(\mathcal{T})$ where $t_0 < \dots < t_k$, then we will write $\alpha := \{t_0 < \dots < t_k\}$ and $\min \alpha := t_0$. $\text{Flag}(\mathcal{T})$ is an abstract simplicial complex with vertex set \mathcal{T} and with k -simplices the elements of $\text{Flag}(\mathcal{T})$ of cardinality $k + 1$. The corresponding topological simplicial complex is the *geometric realization* of the poset \mathcal{T} and is denoted by $|\mathcal{T}|$. For example, if \mathcal{P} is the poset of proper subsets of S , partially ordered by inclusion, then its opposite poset, \mathcal{P}^{op} , is the poset of nonempty faces of the simplex Δ . $\text{Flag}(\mathcal{P})$ is the poset of simplices in its barycentric subdivision, $b\Delta$, and $|\mathcal{P}| = b\Delta$.

The standard realization, $\mathcal{U}(\Phi, K)$. As before, \mathcal{S} denotes the poset of spherical subsets of S . Put $K := |\mathcal{S}|$. It is a subcomplex of $b\Delta$ (provided W is infinite). The mirror structure on Δ induces one on K . More specifically, for each $s \in S$, put $K_s := |\mathcal{S}_{\geq \{s\}}|$ and for each $T \in \mathcal{S}$, $K_T = |\mathcal{S}_{\geq T}|$. (K is the “compact core” of Δ^f .) K is sometimes called the *Davis chamber* of (W, S) and $\mathcal{U}(W, K)$, the *Davis complex*. Alternatively, $\mathcal{U}(\Phi, K)$ is the geometric realization of the poset of spherical residues of Φ (see [8]).

By construction $\mathcal{U}(\Phi, K)$ is locally finite (since Φ is assumed to have finite thickness). It is proved in [8] that $\mathcal{U}(\Phi, K)$ is contractible.

The realization $\mathcal{U}(\Phi, \Delta^f)$. Δ^f and K have the same poset of faces (indexed by \mathcal{S}) and there is a face-preserving deformation retraction $\Delta^f \rightarrow K$. That is to say, Δ^f is a “thickened version” of K . Similarly, $\mathcal{U}(\Phi, \Delta^f)$ is a thickened version of $\mathcal{U}(\Phi, K)$. Like $\mathcal{U}(\Phi, K)$, the space $\mathcal{U}(\Phi, \Delta^f)$ has the advantage of being locally finite; however, the chamber Δ^f is not compact whenever $\Delta^f \neq \Delta$.

3 A CAT(0) metric on $\mathcal{U}(\Phi, \Delta^f)$

Our goal in this section is to prove the following.

Theorem 3.1. *Let Φ be a building of type (W, S) with W infinite. Then there is a piecewise Euclidean, CAT(0) metric on $\mathcal{U}(\Phi, \Delta^f)$.*

Review of the Moussong metric. Suppose T is a spherical subset of S . W_T acts on \mathbf{R}^T via the canonical representation. The *Coxeter cell* of type T , denoted P_T , is defined to be the convex hull of the W_T -orbit of a point x_0 in the interior of the fundamental simplicial cone. As examples, if W_T is a product of n copies of the cyclic group of order 2, then P_T is an n -cube; if W_T is the symmetric group on $n + 1$ letters, then P_T is an n -dimensional permutohedron. Its boundary complex, ∂P_T , is the dual of the Coxeter complex of W_T (the Coxeter complex is a triangulation of the unit sphere in \mathbf{R}^T). The fact that ∂P_T is dual to a simplicial complex means that P_T is a “simple polytope”. The isometry type of P_T is determined once we choose the distance from x_0 to each of the bounding hyperplanes of the simplicial cone. (We assume, without further comment, that such a choice of distance has been made for each $s \in S$.) The intersection of P_T with the fundamental simplicial cone is denoted B_T and called the *Coxeter block* of type T . It is a convex cell combinatorially isomorphic to a cube of dimension $|T|$ (because P_T is simple). One can identify B_T with the subcomplex $|\mathcal{S}_{\leq T}|$ of K in such a way that x_0 is identified with the vertex corresponding to \emptyset . To be more precise, B_T is the union of simplices of $\text{Flag}(\mathcal{S})$ whose maximum vertex is $\leq T$, i.e., $|\mathcal{S}_{\leq T}|$ is a subdivision of B_T .

The convex polytope B_T has two types of faces. First, there are the faces which contain the vertex x_0 . Each such face is a Coxeter block of the form B_V for some $V \subseteq T$. The other type of face is the intersection of B_T with the face of the fundamental simplicial cone fixed by W_V for some nonempty $V \subseteq T$. We denote such a face by $B_{T,V}$ and call it a *reflecting face* of B_T . For the purpose of unifying different cases, we shall sometimes write $B_{T,\emptyset}$ instead of B_T .

The *Moussong metric* on K is the piecewise Euclidean metric on K in which each Coxeter block B_T is given its Euclidean metric as a convex cell in \mathbf{R}^T (cf. [20] or [9, Section 12.1]). This induces a piecewise Euclidean metric on $\mathcal{U}(W, K)$ as well as one on $\mathcal{U}(\Phi, K)$. The link of the central vertex corresponding to \emptyset can be identified with a certain simplicial complex L ($= L(W, S)$) called the *nerve* of the Coxeter system. The vertex set of L is S and a subset $T \subseteq S$ spans a simplex if and only if it is spherical.

Thus, the poset of simplices in L (including the empty simplex) is \mathcal{S} . The piecewise Euclidean metric on K induces a piecewise spherical metric on L such that whenever $m_{st} < \infty$, the length of the edge corresponding to $\{s, t\}$ is $\pi - \pi/m_{st}$. Moussong proved that this piecewise spherical metric on L is CAT(1) and from this he deduced that the piecewise Euclidean metric on $\mathcal{U}(W, K)$ is CAT(0) (cf. [20] and [9, Section 12.3]). Using this, it is proved in [8] that for any building Φ , the Moussong metric on the standard realization, $\mathcal{U}(\Phi, K)$, is CAT(0).

A piecewise Euclidean metric on Δ^f . We will define a cell structure on Δ^f so that each cell will have the form $B_{T,V} \times [0, \infty)^m$ for some $T \in \mathcal{S}$, $V \subseteq T$ and nonnegative integer m . When $m > 0$ such a cell will be noncompact. There are two types of such cells. First there are the compact cells $B_{T,V}$ where $T \in \mathcal{S}$ and $V \subseteq T$. The remaining cells are in bijective correspondence with triples (T, V, α) where $T \in \mathcal{S}$, $V \subseteq T$ and $\alpha \in \text{Flag}(\mathcal{P} - \mathcal{S})$ is such that $T < \min \alpha$. Let \mathcal{F} be the set consisting of pairs (T, V) and triples (T, V, α) , where $T \in \mathcal{S}$, $V \subseteq T$ and $\alpha \in \text{Flag}(\mathcal{P} - \mathcal{S})$ is such that $T < \min \alpha$. \mathcal{F} is partially ordered as follows:

- $(T', V') \leq (T, V)$ if and only if $T' \subseteq T$ and $V' \subseteq V$,
- $(T', V', \alpha') \leq (T, V, \alpha)$ if and only if $(T', V') \leq (T, V)$ and $\alpha' \subseteq \alpha$,
- $(T', V') \leq (T, V, \alpha)$ if and only if $(T', V') \leq (T, V)$.

The cell $c(T, V, \alpha)$ which corresponds to (T, V, α) is defined to be $B_{T,V} \times [0, \infty)^\alpha$, where $[0, \infty)^\alpha$ means the set of all functions from the finite set α to $[0, \infty)$. For the most part, it will suffice to deal with the case $V = \emptyset$ since the cells of the form $B_T \times [0, \infty)^\alpha$ cover Δ^f . The piecewise Euclidean structure on Δ^f will be defined by declaring each $B_T \times [0, \infty)^\alpha$ to have the product metric. Thus, the piecewise Euclidean metric on Δ^f will extend the one on K .

Lemma 3.2. *Δ^f has a decomposition into the cells, $\{B_{T,V}\} \cup \{c(T, V, \alpha)\}$, defined above, where (T, V) and (T, V, α) range over \mathcal{F} .*

To prove this, we need to set up a standard identification of the open cone of radius 1 on a k -simplex σ with the standard simplicial cone $[0, \infty)^{k+1} \subset \mathbf{R}^{k+1}$. Let $\{v_b\}_{b \in \mathcal{B}}$ be the vertex set of σ for some finite index set \mathcal{B} and let $(x_b)_{b \in \mathcal{B}}$ be barycentric coordinates on σ . Let $\mathbf{R}^\mathcal{B}$ be the Euclidean space of all functions $\mathcal{B} \rightarrow \mathbf{R}$. Let $\mathbb{S}_+(\mathbf{R}^\mathcal{B})$ denote the intersection of the standard simplicial cone $[0, \infty)^\mathcal{B}$ with the unit sphere. Thus, $\mathbb{S}_+(\mathbf{R}^\mathcal{B})$ is an ‘‘all right’’ spherical simplex (i.e., all edge lengths and all dihedral angles are $\pi/2$). Let

$\{e_b\}_{b \in \mathcal{B}}$ be the standard basis for $\mathbf{R}^{\mathcal{B}}$. Define a homeomorphism $\theta_{\mathcal{B}} : \sigma \rightarrow \mathbb{S}_+(\mathbf{R}^{\mathcal{B}})$, taking v_b to e_b , by

$$\sum x_b v_b \rightarrow \sum x_b e_b \Big/ \left(\sum x_b^2 \right)^{1/2}. \quad (3.1)$$

Let $r : [0, 1) \rightarrow [0, \infty)$ be some fixed homeomorphism. The open cone on σ can be regarded as the points $[t, x]$ in the join, $v * \sigma$, of σ with a point v such that the join coordinate t is $\neq 1$. The homeomorphism from the open cone to $[0, \infty)^{\mathcal{B}}$ is defined by $[t, x] \rightarrow r(t)\theta_{\mathcal{B}}(x)$.

Proof of Lemma 3.2. Suppose $\alpha \in \text{Flag}(\mathcal{P})$. Then $\alpha = \{T_0 < \dots < T_{k+l}\}$, where $T_k \in \mathcal{S}$ and $T_{k+1} \notin \mathcal{S}$. Put $\alpha' := \{T_0 < \dots < T_k\}$ and $\alpha'' := \{T_{k+1} < \dots < T_{k+l}\}$. The simplex $\sigma_{\alpha'}$ lies in B_{T_k} while the simplex $\sigma_{\alpha''}$ is in the nonspherical face $\Delta_{T_{k+1}}$. We have $\sigma_{\alpha} = \sigma_{\alpha'} * \sigma_{\alpha''}$ and a point in σ_{α} has coordinates $[t, x, y]$ where $t \in [0, 1]$, $x \in \sigma_{\alpha'}$ and $y \in \sigma_{\alpha''}$. The points in $\sigma_{\alpha} - \sigma_{\alpha''}$ are those where the join coordinate t is $\neq 1$. The identification $\sigma_{\alpha} - \sigma_{\alpha''} \rightarrow \sigma_{\alpha'} \times [0, \infty)^{\alpha''}$ is given by $[t, x, y] \rightarrow (x, r(t)\theta_{\alpha''}(y))$. \square

Next we want to consider the link of the central vertex v_{\emptyset} (corresponding to \emptyset) in this cell structure. Let Δ^{op} denote the simplex on S . The nerve L is a subcomplex of Δ^{op} . If W is spherical, then $L = \Delta^{op}$, while if W is infinite, then L is a subcomplex of $\partial\Delta^{op}$. Moreover, $\partial\Delta^{op}$ is a triangulation of the $(n-1)$ -sphere, for $n = |S|$. Assume W is infinite. The link of v_{\emptyset} in Δ^f is a certain subdivision L' of $\partial\Delta^{op}$, which we shall now describe. The vertex set of L' is the disjoint union $S \cup (\mathcal{P} - \mathcal{S})$. There are three types of simplices in L' :

- 1) simplices σ_T in L corresponding to spherical subsets $T \in \mathcal{S}_{>\emptyset}$,
- 2) simplices σ_{α} in $b\partial\Delta (= b\partial\Delta^{op})$ corresponding to flags $\alpha \in \text{Flag}(\mathcal{P} - \mathcal{S})$,
- 3) joins $\sigma_T * \sigma_{\alpha}$, with $T \in \mathcal{S}_{>\emptyset}$, $\alpha \in \text{Flag}(\mathcal{P} - \mathcal{S})$ and $T < \min \alpha$.

Lemma 3.3. *L' is a subdivision of $\partial\Delta^{op}$ and $L \subseteq L'$ is a full subcomplex.*

Proof. Suppose U is a minimal element of $\mathcal{P} - \mathcal{S}$. Let σ_U denote the corresponding simplex in $\partial\Delta^{op}$. Introduce a ‘‘barycenter’’ $v_U \in \partial\Delta^{op}$ and then subdivide σ_U to a new simplicial complex $(\sigma_U)'$ by coning off the simplices in $\partial\sigma_U$. Each new simplex will have the form $v_U * \sigma_T$ for some $T \subset U$.

Next, let U be an arbitrary element of $\mathcal{P} - \mathcal{S}$ and suppose by induction that we have defined the subdivision of $(\sigma_{U'})$ for each $U' \subset U$ and hence, a subdivision $(\partial\sigma_U)'$ of $\partial\sigma_U$. Introduce a barycenter v_U of σ_U and subdivide

by coning off $(\partial\sigma_U)'$. Each new simplex will have the form $v_U * \sigma_{U'}$ for some $U' \subset U$. In other words, each new simplex will be either of type 2 or type 3 above. It is clear that the subcomplex L is full. \square

The following lemma is also clear.

Lemma 3.4. *L' is the link of the central vertex in the piecewise Euclidean cell structure on Δ^f .*

The piecewise Euclidean metric induces a piecewise spherical metric on L' extending the given metric on L . Since the link of the origin in $[0, \infty)^{k+1}$ is the all right spherical k -simplex we get the following description of the metric on L' .

Lemma 3.5. *The simplices in L' have spherical metrics of the following types.*

- 1) *For $T \in \mathcal{S}_{>0}$, the simplex σ_T in L is the dual to the fundamental simplex for W_T on the unit sphere in \mathbf{R}^T . In other words, for distinct elements s, t in T , the edge corresponding to $\{s, t\}$ has length $\pi - \pi/m_{st}$.*
- 2) *The simplex σ_α corresponding to $\alpha \in \text{Flag}(\mathcal{P} - \mathcal{S})$ has its all right structure. In other words, each edge of σ_α has length $\pi/2$.*
- 3) *The simplex $\sigma_T * \sigma_\alpha$ has the structure of a spherical join. In other words, the length of an edge connecting a vertex in σ_T to one in σ_α is $\pi/2$.*

Remark 3.6. The simplicial complex L' is the nerve of a Coxeter group (W', S') which contains W as a special subgroup. Namely, S' is the disjoint union, $S \cup (\mathcal{P} - \mathcal{S})$. Two generators of S are related as before. If $U, V \in \mathcal{P} - \mathcal{S}$, then put $m(U, V) := 2$ whenever $U \subset V$ or $V \subset U$ and $m(U, V) := \infty$, otherwise. Similarly, if $s \in S$ and $U \in \mathcal{P} - \mathcal{S}$, then $m(s, U) = m(U, s) = 2$ when $s \in U$ and it is $= \infty$, otherwise. Since L' is a triangulation of S^{n-1} , with $n = |S|$, this shows that any n generator Coxeter group is a special subgroup of a Coxeter group which acts cocompactly on a contractible n -manifold (such a Coxeter group is said to be *type HM^n* in [9]).

CAT(0) and CAT(1) metrics. Gromov [18] proved that a piecewise Euclidean metric on a polyhedron Y is locally CAT(0) if and only if the link in Y of each cell is CAT(1). In his proof that $L(W, S)$ was CAT(1), Moussong [20] gave a criteria for certain piecewise spherical structures on simplicial complexes to be CAT(1). We recall his criteria below. A spherical simplex *has size $\geq \pi/2$* if each of its edges has length $\geq \pi/2$. A spherical

simplex $\sigma \subset \mathbb{S}^k \subset \mathbf{R}^{k+1}$ with vertex set $\{v_0, \dots, v_{k+1}\}$ and edge lengths $l_{ij} := \cos^{-1}(v_i \cdot v_j)$ is determined up to isometry by the (l_{ij}) . Conversely, a symmetric $(k+1) \times (k+1)$ matrix (l_{ij}) of real numbers in $[0, \pi)$ can be realized as the set of edge lengths of a spherical simplex if and only if the matrix $(\cos l_{ij})$ is positive definite, cf. [9, Lemma I.5.1, p. 513]. Suppose N is a simplicial complex with a piecewise spherical structure (i.e., each simplex has the structure of a spherical simplex). N has size $\geq \pi/2$ if each of its simplices does. N is a *metric flag complex* if it satisfies the following condition: given any collection of vertices $\{v_0, \dots, v_k\}$ which are pairwise connected by edges, then $\{v_0, \dots, v_k\}$ is the vertex set of a simplex in N if and only if the matrix of edge lengths (l_{ij}) can be realized as the matrix of edge lengths of an actual spherical simplex. (In other words, if and only if $(\cos l_{ij})$ is positive definite.) Moussong's Lemma is the following.

Lemma 3.7. (Moussong's Lemma, [20] or [9, Appendix I.7]). *Suppose a piecewise spherical simplicial complex N has size $\geq \pi/2$. Then N is CAT(1) if and only if it is a metric flag complex.*

The next result follows immediately from our previous description of the piecewise spherical complex L' .

Lemma 3.8. *L' has size $\geq \pi/2$ and is a metric flag complex.*

Corollary 3.9. *L' is CAT(1). Moreover, L is a totally geodesic subcomplex.*

Proof. The last sentence follows from the fact that L is a full subcomplex. \square

Since the link of any simplex in a metric flag complex of size $\geq \pi/2$ has the same properties (cf. [20, Lemma 8.3] or [9, Lemma I.5.11]), the link of any simplex in L' is also CAT(1). Since the link in $\Delta^f \cap \Delta_V$ of any cell of the form $B_{T,V} \times [0, \infty)^\alpha$ can be identified with the link of the corresponding simplex $\sigma_T * \sigma_\alpha$ in L' , it follows that the link of each cell in Δ^f is CAT(1).

Corollary 3.10. *The piecewise Euclidean metric on $\mathcal{U}(W, \Delta^f)$ is CAT(0).*

Proof. The union of W_T -translates of a Coxeter block B_T in $\mathcal{U}(W, \Delta^f)$ is a Coxeter cell P_T and the complete inverse image of B_T in $\mathcal{U}(W, \Delta^f)$ is a disjoint union of copies of P_T . Hence, $\mathcal{U}(W, \Delta^f)$ has a cell structure in which the cells are either translates of Coxeter cells of the form P_T or translates of cells of the form $P_T \times [0, \infty)^\alpha$ for some $\alpha \in \text{Flag}((\mathcal{P} - \mathcal{S})_{>T})$. In either case the link of such a cell in $\mathcal{U}(W, \Delta^f)$ is identified with the link of the corresponding cell in Δ^f . By Corollary 3.9, $\mathcal{U}(W, \Delta^f)$ is locally CAT(0). A

space is CAT(0) if and only if it is locally CAT(0) and simply connected (cf. [18, p. 119] or [5, Ch. II.4]). $\mathcal{U}(W, \Delta^f)$ is contractible (hence, simply connected), since it is homotopy equivalent to $\mathcal{U}(W, K)$. Therefore, it is CAT(0). \square

We also want to consider links of cells of the form $B_{T,V}$ or $B_{T,V} \times [0, \infty)^\alpha$ in various cell complexes. (Here $B_{T,V}$ is a reflecting face of B_T .)

Lemma 3.11. *Suppose c is a cell in Δ^f of the form $c = B_{T,V}$ or $c = B_{T,V} \times [0, \infty)^\alpha$ for some $V \subseteq T \in \mathcal{S}_{>\emptyset}$ and $\alpha \in \text{Flag}((\mathcal{P} - \mathcal{S})_{>T})$. Let $d = B_T$ or $B_T \times [0, \infty)^\alpha$ be the corresponding larger cell in Δ^f and let σ_d be the corresponding simplex in L' . Then*

$$\text{Lk}(c, \Delta^f) = \text{Lk}(\sigma_d, L') * \tau(V) \quad (3.2)$$

$$\text{Lk}(c, \mathcal{U}(W, \Delta^f)) = \text{Lk}(\sigma_d, L') * \mathbb{S}^V \quad (3.3)$$

Here \mathbb{S}^V is the unit sphere in the canonical representation of W_V on \mathbf{R}^V and $\tau(V) \subset \mathbb{S}^V$ is the fundamental simplex.

Moreover, suppose that Φ is a building of type (W, S) , that R is the spherical residue of type V containing the base chamber and that $\mathbb{S}(R) := \mathcal{U}(R, \tau(V))$ is the spherical realization of R . Then

$$\text{Lk}(c, \mathcal{U}(\Phi, \Delta^f)) = \text{Lk}(\sigma_d, L') * \mathbb{S}(R). \quad (3.4)$$

Proof. Let Δ_V^f denote the face of Δ^f fixed by W_V (cf. (2.1)). Then

$$\begin{aligned} \text{Lk}(c, \Delta^f) &= \text{Lk}(c, \Delta_V^f) * \text{Lk}(\Delta_V^f, \Delta^f) \\ &= \text{Lk}(d, \Delta^f) * \tau(V) \\ &= \text{Lk}(\sigma_d, L') * \tau(V) \end{aligned}$$

and similarly for formulas (3.3) and (3.4). \square

We can now prove the main result of this section.

Proof of Theorem 3.1. Any spherical building, such as $\mathbb{S}(R)$, is CAT(1) (e.g., see [8]) and the spherical join of two CAT(1)-spaces is CAT(1) (e.g., see [5]). So, the theorem follows from (3.4). Alternatively, it can be proved from Corollary 3.10 by using the argument in [8, §11]. \square

A variation. In Section 4 we will need the following modification of the previous construction. Given a subset $U \subseteq S$, we will define a new piecewise

Euclidean metric on $\Delta^f - \Delta^U$ and then show that it induces CAT(0) metrics on $\mathcal{U}(W_{S-U}, \Delta^f - \Delta^U)$ and $\mathcal{U}(R, \Delta^f - \Delta^U)$ for any $(S-U)$ -residue R of Φ .

For each spherical subset T , let $C_T^* \subset \mathbf{R}^T$ be the simplicial cone determined by the bounding hyperplanes of B_T passing through the vertex x_0 . (In other words, C_T^* is the dual cone to the fundamental simplicial cone.) Let $\mathcal{S}_{S-U} := \mathcal{S}(W_{S-U}, S-U)$ be the poset of spherical subsets of $S-U$ and let $L_{S-U} := L(W_{S-U}, S-U)$ be the nerve of W_{S-U} . $\Delta^f - \Delta^U$ has a cell structure with cells of the following two types:

- (a) $B_{T,V}$, where $T \in \mathcal{S}_{S-U}$ and $V \subseteq T$,
- (b) $B_T \times [0, \infty)^\alpha$, where $\alpha \in \mathcal{P} - \mathcal{S}_{S-U}$ and T, V are as above.

As before, each such cell is given the natural product metric. In effect we are putting $\Upsilon(\Delta) \cup \Delta^U$ at infinity.

The piecewise Euclidean metric on $\Delta^f - \Delta^U$ induces one $\mathcal{U}(R, \Delta^f - \Delta^U)$. Let us describe the link, L'_{S-U} , of the central vertex in the new metric on $\Delta^f - \Delta^U$. The vertex set of L'_{S-U} is $(S-U) \cup (\mathcal{P} - \mathcal{S}_{S-U})$. As before, there are three types of simplices:

- 1) simplices σ_T in L_{S-U} corresponding to spherical subsets $T \in \mathcal{S}_{S-U}$,
- 2) simplices σ_α in $b\partial\Delta$ ($= b\partial\Delta^{op}$) corresponding to flags $\alpha \in \text{Flag}(\mathcal{P} - \mathcal{S}_{S-U})$,
- 3) joins $\sigma_T * \sigma_\alpha$, with $T \in \mathcal{S}_{S-U}$, $\alpha \in \text{Flag}(\mathcal{P} - \mathcal{S}_{S-U})$ and $T < \min \alpha$.

and L'_{S-U} can be identified with a subdivision of $\partial\Delta^{op}$. Moreover, just as before, L'_{S-U} has size $\geq \pi/2$ and is a metric flag complex; hence, it is CAT(1). This proves the following.

Theorem 3.12. *Suppose $U \subseteq S$ and R is any $(S-U)$ -residue in Φ . Then the piecewise Euclidean metric on $\mathcal{U}(R, \Delta^f - \Delta^U)$, defined above, is CAT(0).*

4 Metric spheres in $\mathcal{U}(\Phi, \Delta^f)$

An n -dimensional cell complex X is *CM* (for ‘‘Cohen-Macaulay’’) if $\tilde{H}^*(X)$ is concentrated in degree n and is a free abelian group in that degree. Similarly, an n -dimensional, noncompact, contractible space X is *SI* (for ‘‘spherical at infinity’’) if $H_c^*(X)$ is concentrated in degree n and is a free abelian group in that degree. We want to prove that $\mathcal{U}(\Phi, \Delta^f)$ is *SI*. This is a consequence of Theorems 4.1 and 4.2 below.

Suppose N is a CAT(1), piecewise spherical polyhedron and that $p \in N$. Let $B(p, \pi/2) \subseteq N$ denote the open ball of radius $\pi/2$ centered at p . Define a space PN_p , called N punctured at p , by $PN_p := N - B(p, \pi/2)$. We are interested in this concept when $N = \text{Lk}(c)$, the link of a cell c in some CAT(0) complex X . In this case we will write $\text{PLk}_p(c)$ for PN_p and call it the *punctured link* of c at p .

Theorem 4.1. (Brady, McCammond and Meier [4]). *Let X be a CAT(0), piecewise Euclidean cell complex (with finitely many shapes of cells). If for each cell c in X and for each $p \in \text{Lk}(c)$, the spaces $\text{Lk}(c)$ and $\text{PLk}_p(c)$ are $(n - \dim c)$ -acyclic, then X is n -acyclic at infinity. In particular, if X satisfies this condition and is n -dimensional, then it is SI.*

Remarks. In [4] the hypothesis of the above theorem is that X is the universal cover of a finite, nonpositively curved complex; however, the proof clearly works with a weaker hypothesis such as finitely many shapes of cells (which holds in our case). The proof of the theorem uses Morse theory for polyhedral complexes. Roughly, it goes as follows. Let $\rho : X \rightarrow \mathbf{R}$ be the distance from some base point x_0 , i.e., $\rho(x) := d(x, x_0)$. The spheres $S(r)$ of radius r centered at $x_0 \in X$ are the level sets of ρ . Call a point x a *critical point* (of ρ) if it is the closest point to x_0 in some closed cell c . It follows from the CAT(0) hypothesis that the critical points are isolated. If, for sufficiently small ε , there is no critical point in the annular region between $S(r + \varepsilon)$ and $S(r)$, then $S(r + \varepsilon)$ and $S(r)$ are homeomorphic. On the other hand, the effect of crossing a critical point $x \in S(r)$ is to remove a contractible neighborhood of x in $S(r)$ and replace it by the punctured link $\text{PLk}(x)_p$ where p is the direction at x of the geodesic from x to x_0 . (If x lies in the relative interior of a k -dimensional cell c , then $\text{Lk}(x) \cong \mathbb{S}^{k-1} * \text{Lk}(c)$.) So, the effect on the homotopy type of the level sets is to replace a contractible neighborhood by a copy of a suspension of a punctured link. It follows that, under the hypotheses of Theorem 4.1, each metric sphere is CM. Since X is CAT(0), metric balls are contractible and since $H_c^*(X) = \varinjlim H^*(B(r), S(r))$ is concentrated in the top degree, X is SI.

Theorem 4.2. (Dymara–Osajda [16] and Schulz [24]). *Suppose R is a spherical building of type (W_T, T) and that $\mathbb{S}(R)$ ($:= \mathcal{U}(R, \tau(T))$) is its spherical realization. Then for any $p \in \mathbb{S}(R)$, the space $\text{PS}(R)_p$ is CM.*

An immediate corollary to Theorems 4.1 and 4.2 is the following.

Theorem 4.3. (cf. [2]). *With notation as in Section 3 and above, given any building Φ , its realization $\mathcal{U}(\Phi, \Delta^f)$ is SI.*

As a corollary to Theorem 3.12 we get the following relative version of Theorem 4.1.

Theorem 4.4. *Suppose $U \subseteq S$ and R is any $(S - U)$ -residue in Φ . Then $\mathcal{U}(R, \Delta^f - \Delta^U)$ is SI.*

Corollary 4.5. *Suppose $U \subseteq S$. Then $\mathcal{U}(\Phi, \Delta^f - \Delta^U)$ is SI.*

Proof. $\mathcal{U}(\Phi, \Delta^f - \Delta^U)$ is the disjoint union of the spaces $\mathcal{U}(R, \Delta^f - \Delta^U)$ where R ranges over the $(S - U)$ -residues. By the previous theorem, the compactly supported cohomology of each such component is free abelian and concentrated in the top degree. \square

5 Cohomology with finite support and compact support

Given a CW complex Y , $C_{\text{fin}}^*(Y)$ denotes the complex of finitely supported cellular cochains on Y and $H_{\text{fin}}^*(Y)$ its cohomology. When Y is only required to be a topological space, $H_c^*(Y)$ denotes its compactly supported singular cohomology, i.e.,

$$H_c^*(Y) := \varinjlim H^*(Y, Y - C),$$

where the direct limit is over all compact subsets $C \subseteq Y$. If Y is a locally finite CW complex, $H_{\text{fin}}^*(Y) \cong H_c^*(Y)$.

Suppose Z, Z' are mirrored spaces over S . A map $F : Z \rightarrow Z'$ is *mirrored* if $F(Z_T) \subseteq Z'_T$ for all $T \subseteq S$; it is a *mirrored homotopy equivalence* if $F|_{Z_T} : Z_T \rightarrow Z'_T$ is a homotopy equivalence for all T (including $T = \emptyset$).

Given a mirrored space X , let us say that $\Upsilon(X)$ is *collared in X* if there is an increasing family $\mathcal{N} = \{N_\varepsilon\}_{\varepsilon \in (0, a]}$ of open neighborhoods of $\Upsilon(X)$ (“increasing means that $N_\varepsilon \subseteq N_{\varepsilon'}$ whenever $\varepsilon < \varepsilon'$ ”) such that the following two properties hold:

- (i) $\bigcap_{\varepsilon \in (0, a]} N_\varepsilon = \Upsilon(X)$ and
- (ii) For each $\varepsilon > 0$, the inclusion $\Upsilon(X) \hookrightarrow \overline{N_\varepsilon}$ is a mirrored homotopy equivalence.

For example, if X is the simplex Δ , then $\Upsilon(\Delta)$ is collared in Δ . (Proof: $\Upsilon(\Delta)$ is a union of closed faces and we can take N_ε to be its ε -neighborhood in Δ .) More generally, if X is any finite CW complex, then $\Upsilon(X)$ is collared in X .

Property (ii) implies that $\mathcal{U}(\Phi, \Upsilon(X)) \hookrightarrow \mathcal{U}(\Phi, N_\varepsilon)$ is a homotopy equivalence. To simplify notation, in what follows we often write \mathcal{U}_X instead of $\mathcal{U}(\Phi, X)$.

Lemma 5.1. *Suppose X is a finite, mirrored CW complex. Then*

$$H_c^*(\mathcal{U}_{X^f}) = H_{\text{fin}}^*(\mathcal{U}_X, \mathcal{U}_{\Upsilon(X)}).$$

Proof. Since X is a finite complex, $H_{\text{fin}}^*(\mathcal{U}_X, \mathcal{U}_{\Upsilon(X)}) = H_c^*(\mathcal{U}_X, \mathcal{U}_{\Upsilon(X)})$. Since $\Upsilon(X)$ is collared in X , we have a family $\mathcal{N} = \{N_\varepsilon\}$ of open neighborhoods. For a given $N = N_\varepsilon \in \mathcal{N}$, let ∂N denote the boundary of \overline{N} . Since N is mirrored homotopy equivalent to $\Upsilon(X)$,

$$H_{\text{fin}}^*(\mathcal{U}_X, \mathcal{U}_{\Upsilon(X)}) \cong H_c^*(\mathcal{U}_X, \mathcal{U}_{\overline{N}}) \cong H_c^*(\mathcal{U}_{(X-N)}, \mathcal{U}_{\partial N}),$$

where the second isomorphism is an excision. As $\varepsilon \rightarrow 0$, $X - N_\varepsilon \rightarrow X^f$, so any compact subset C of \mathcal{U}_{X^f} lies within some $\mathcal{U}_{X-N_\varepsilon}$. Hence,

$$H_c^*(\mathcal{U}_{X^f}) = \varinjlim H^*(\mathcal{U}_{X^f}, \mathcal{U}_{X^f} - C) = H_c^*(\mathcal{U}_{X-N}, \mathcal{U}_{\partial N}).$$

Combining these equations, we get the result. \square

Example 5.2. Δ^f is a thickened version of K and $\mathcal{U}(\Phi, \Delta^f)$ is a thickened version of $\mathcal{U}(\Phi, K)$. Hence, $\mathcal{U}(\Phi, \Delta^f)$ and $\mathcal{U}(\Phi, K)$ are homotopy equivalent. On the other hand, their compactly supported cohomology groups can be completely different. For example, suppose W is the free product of three copies of $\mathbf{Z}/2$. Then Δ^f is a triangle with its vertices deleted while K is a tripod. $\mathcal{U}(W, \Delta^f)$ can be identified with the hyperbolic plane while $\mathcal{U}(W, K)$ is the regular trivalent tree. (This is the familiar picture of the congruence 2 subgroup of $PSL(2, \mathbf{Z})$ acting on the hyperbolic plane.) The compactly supported cohomology of $\mathcal{U}(W, \Delta^f)$ is that of the plane (i.e., it is concentrated in degree 2 and is isomorphic to \mathbf{Z} in that degree), while the compactly supported cohomology of $\mathcal{U}(W, K)$ is that of a tree (i.e., it is concentrated in degree 1 and is a countably generated, free abelian group in that degree).

6 Cohomology with coefficients in $\mathcal{I}(A)$

Let $A (= A(\Phi))$ be the free abelian group of finitely supported, \mathbf{Z} -valued functions on Φ . For each subset T of S , define a \mathbf{Z} -submodule of A :

$$A^T := \{f \in A \mid f \text{ is constant on each residue of type } T\}.$$

Note that $A^\emptyset = A$. Also note that $A^T = 0$ whenever T is not spherical.

We have $A^U \subset A^T$ when $T \subset U$. Let $A^{>T}$ be the \mathbf{Z} -submodule of A^T spanned by the A^U with T a proper subset of U . Put

$$D^T := A^T / A^{>T}.$$

Remark 6.1. When $\Phi = W$, A is the group ring $\mathbf{Z}W$ and A^T consists of elements in $\mathbf{Z}W$ which are constant on each left coset wW_T . Let us assume T is spherical (otherwise $A^T = 0$). Let $\text{In}(w) := \{s \in S \mid l(ws) < l(w)\}$ be the set of letters with which a reduced expression for w can end. Since T is spherical, each left coset of W_T has a unique representative w which has a reduced expression ending in the longest element of W_T ; hence, this representative has $T \subseteq \text{In}(w)$. As a basis for D^T , we can take images of the elements in A^T corresponding to cosets which have longest representatives with $T = \text{In}(w)$.

Definition 6.2. The abelian group A and the family of subgroups $\{A^T\}_{T \subseteq S}$ define a ‘‘coefficient system’’ $\mathcal{I}(A)$ on X so that the *i-cochains with coefficients in $\mathcal{I}(A)$* are given by

$$\mathcal{C}^i(X; \mathcal{I}(A)) := \prod_{c \in X^{(i)}} A^{S(c)}$$

where $X^{(i)}$ denotes the set of i -cells in X .

We continue to write \mathcal{U}_X for $\mathcal{U}(\Phi, X)$. Let $\mathcal{U}^{(i)}$ denote the set of i -cells in \mathcal{U}_X . Given a chamber $\varphi \in \Phi$ and an i -cell c in X , let $\varphi \cdot c$ denote the corresponding i -cell in \mathcal{U}_X . Given a finitely supported function $\alpha : \mathcal{U}^{(i)} \rightarrow \mathbf{Z}$, and an i -cell $c \in X^{(i)}$ with $S(c)$ spherical, we get an element $f \in A^{S(c)}$ defined by $f(\varphi) := \alpha(\varphi \cdot c)$. Of course, when $S(c)$ is not spherical, $A^{S(c)} = 0$. So, for any finite, mirrored CW complex X , this establishes an isomorphism

$$\mathcal{C}^i(X; \mathcal{I}(A)) \cong C_{\text{fin}}^i(\mathcal{U}_X, \mathcal{U}_{\Gamma(X)}). \quad (6.1)$$

In other words, the isomorphism (6.1) is given by identifying a finitely supported function on the inverse image in $\mathcal{U}^{(i)}$ of a cell $c \in X^{(i)}$ with a function on Φ (i.e., with an element of A) which is constant on $S(c)$ -residues containing the cells $\varphi \cdot c$.

The coboundary maps in $\mathcal{C}^*(X; \mathcal{I}(A))$ are defined by using these isomorphisms to transport the coboundary maps on finitely supported cochains to $\mathcal{C}^*(X; \mathcal{I}(A))$. This means that the coboundary maps in $\mathcal{C}^*(X; \mathcal{I}(A))$ are defined by combining the usual coboundary maps in $C^*(X)$ with the inclusions $A^U \hookrightarrow A^T$ for $U \supset T$. So, we have the following.

Lemma 6.3. *For X a finite, mirrored CW complex,*

$$\mathcal{H}^*(X; \mathcal{I}(A)) = H_{\text{fin}}^*(\mathcal{U}_X, \mathcal{U}_{\Upsilon(X)}) = H_c^*(\mathcal{U}_{X^f}).$$

Proof. The second equation is from Lemma 5.1. \square

Remark 6.4. If $\Phi = W$, then $A = \mathbf{Z}W$. $\mathcal{C}^*(X; \mathcal{I}(\mathbf{Z}W))$ can be interpreted as the equivariant cochains on $\mathcal{U}(W, X)$ with coefficients in $\mathbf{Z}W$. When X is a finite complex, Lemma 6.3 asserts

$$\mathcal{C}^*(X; \mathcal{I}(\mathbf{Z}W)) = C_{\text{fin}}^*(\mathcal{U}(W, X), \mathcal{U}(W, \Upsilon(X))).$$

The corresponding cohomology groups were studied in [10, 7]. In particular, when $X = K$, these cohomology groups are isomorphic to $H^*(W; \mathbf{Z}W)$.

In what follows the coefficient system is usually $\mathcal{I}(A)$ and we shall generally omit it from our notation, writing $\mathcal{H}(\)$ and $\mathcal{C}(\)$ instead of $\mathcal{H}(\ ; \mathcal{I}(A))$ and $\mathcal{C}(\ ; \mathcal{I}(A))$. (In other words, the coefficients $\mathcal{I}(A)$ are implicit when we use the calligraphic \mathcal{C} or \mathcal{H} notation.) As usual, Δ is the simplex of dimension $n = |S| - 1$ with its codimension one faces indexed by S .

Theorem 6.5. (cf. [15, Theorem B] as well as [11, 14]). *$\mathcal{H}^*(\Delta)$ is concentrated in degree n and is free abelian; moreover, $\mathcal{H}^n(\Delta) = D^\emptyset$. More generally, for any subset U of S , $\mathcal{H}^*(\Delta, \Delta^U)$ is concentrated in degree n and is free abelian and*

$$\mathcal{H}^n(\Delta, \Delta^U) = A / \sum_{s \in S-U} A^s$$

Proof. By Lemma 6.3, $\mathcal{H}^*(\Delta) = H_c^*(\mathcal{U}_{\Delta^f})$ and by Theorem 4.3, the right hand side is concentrated in degree n and is free abelian. The cochain complex looks like

$$\dots \rightarrow \mathcal{C}^{n-1}(\Delta) \rightarrow \mathcal{C}^n(\Delta) \rightarrow 0,$$

where $\mathcal{C}^n(\Delta) = A$ and $\mathcal{C}^{n-1}(\Delta) = \bigoplus A^s$. It follows that cohomology in degree n is the quotient

$$A / \sum_{s \in S} A^s = D^\emptyset.$$

Similarly, $\mathcal{H}^*(\Delta, \Delta^U) = H_c^*(\mathcal{U}_{\Delta^f - \Delta^U})$, and by Corollary 4.5 the right hand side is concentrated in degree n and is free abelian. Since $\mathcal{C}^n(\Delta, \Delta^U) = A$ and $\mathcal{C}^{n-1}(\Delta, \Delta^U) = \bigoplus_{s \notin U} A^s$, we get the final formula in the theorem. \square

Corollary 6.6. *D^\emptyset is free abelian.*

Remark 6.7. We saw in Section 2 that whenever W is infinite, $\mathcal{U}(W, \Delta^f)$ is homeomorphic to Euclidean space \mathbf{R}^n . Hence, the compactly supported cohomology of $\mathcal{U}(W, \Delta^f)$ is that of $H_c^*(\mathbf{R}^n)$. Similarly, $\mathcal{U}(R, \Delta^f - (\Delta^f)^U)$ is homeomorphic to \mathbf{R}^n , for each $(S - U)$ -residue R .

Remark 6.8. $\mathcal{U}(W, \Delta^f)$ is a thickened version of $\mathcal{U}(W, K)$. In the proof of Corollary 3.10 we explained the cellulation of $\mathcal{U}(W, K)$ by ‘‘Coxeter cells.’’ The corresponding cellular chain complex, $C_*(\mathcal{U}(W, K))$, has the form

$$\mathbf{Z}W \longleftarrow \bigoplus_{s \in S} H^s \longleftarrow \bigoplus_{T \in \mathcal{S}^{(2)}} H^T \longleftarrow \dots$$

where H^T is the representation induced from the sign representation of W_T and where $\mathcal{S}^{(k)}$ is the set of spherical subsets with k elements (cf. [11, §8]). The cochain complex $C^*(\Delta)$ for $n - 2 \leq * \leq n$, looks like

$$\dots \longrightarrow \bigoplus_{T \in \mathcal{S}^{(2)}} A^T \longrightarrow \bigoplus_{s \in S} A^s \longrightarrow \mathbf{Z}W.$$

So, $C_*(\mathcal{U}(W, K))$ and $C_c^{n-*}(\mathcal{U}(W, \Delta^f))$ are Poincaré dual.

7 The Decomposition Theorem

We want to prove a version of Theorem 6.5 for lower dimensional spherical faces of Δ . Suppose T is a spherical subset of S . Put $\sigma = \Delta_T$ and $m = n - |T|$. Let U be an arbitrary subset of $S - T$. We continue the policy of omitting the coefficient system $\mathcal{I}(A)$ from our notation.

Proposition 7.1. *Each of the following cohomology groups is concentrated in the top degree and is a free abelian group in that degree:*

$$\mathcal{H}^*(\sigma, \sigma^U), \mathcal{H}^*(\sigma^U, \partial(\sigma^U)), \text{ and } \mathcal{H}^*(\sigma^U)$$

(The top degrees are m , $m - 1$, and $m - 1$, respectively.) Moreover,

$$\mathcal{H}^m(\sigma, \sigma^U) = A^T / \sum_{s \in (S-T)-U} A^{T \cup \{s\}}, \quad (7.1)$$

$$\mathcal{H}^{m-1}(\sigma^U, \partial(\sigma^U)) = \sum_{s \in U} A^{T \cup \{s\}}, \quad (7.2)$$

$$\mathcal{H}^{m-1}(\sigma^U) = \sum_{s \in U} A^{T \cup \{s\}} / \sum_{\substack{s \in U \\ t \in (S-T)-U}} A^{T \cup \{s, t\}}, \quad (7.3)$$

Proof. We first prove $\mathcal{H}^*(\sigma, \sigma^U)$ is concentrated in degree m and is free abelian. The proof is by induction on the number of elements in T . It holds for $|T| = 0$ by Theorem 6.5. Suppose $T = T' \cup \{s\}$, $U' = U \cup \{s\}$ and $\tau = \Delta_{T'}$. The exact sequence of the triple $(\tau, \tau^{U'}, \tau^U)$ gives

$$\dots \rightarrow \mathcal{H}^{*-1}(\sigma, \sigma^U) \rightarrow \mathcal{H}^*(\tau, \tau^{U'}) \rightarrow \mathcal{H}^*(\tau, \tau^U) \rightarrow \dots$$

(This uses the excision, $\mathcal{H}^{*-1}(\tau^{U'}, \tau^U) \cong \mathcal{H}^{*-1}(\sigma, \sigma^U)$.) By inductive hypothesis, the last two terms are free abelian and concentrated in degree $m+1$. Hence, $\mathcal{H}^*(\sigma, \sigma^U)$ is concentrated in degree m . It is free abelian since it injects into a free abelian group.

That the other cohomology groups are free abelian and are concentrated in the top degree follows from various exact sequences. For example, for $\mathcal{H}^*(\sigma^U, \partial(\sigma^U))$, consider the sequence of the triple $(\sigma, \partial\sigma, \sigma^{S-U})$:

$$\rightarrow \mathcal{H}^{*-1}(\sigma^U, \partial(\sigma^U)) \rightarrow \mathcal{H}^*(\sigma, \partial\sigma) \rightarrow \mathcal{H}^*(\sigma, \sigma^{S-U}) \rightarrow \quad (7.4)$$

where we have used the excision $\mathcal{H}^{*-1}(\partial\sigma, \sigma^{S-U}) \cong \mathcal{H}^{*-1}(\sigma^U, \partial(\sigma^U))$ to identify the first term. The second and third terms are free abelian and concentrated in degree m ; hence, the first term is free abelian and concentrated in degree $m-1$. For $\mathcal{H}^*(\sigma^U)$, we have the exact sequence of the pair (σ, σ^U) :

$$\rightarrow \mathcal{H}^{*-1}(\sigma^U) \rightarrow \mathcal{H}^*(\sigma, \sigma^U) \rightarrow \mathcal{H}^*(\sigma) \rightarrow \quad (7.5)$$

where the second and third terms are free abelian and concentrated in degree m . Hence, $\mathcal{H}^*(\sigma^U)$ is free abelian and is concentrated in degree $m-1$.

It remains to verify formulas (7.1), (7.2) and (7.3). We have $\mathcal{C}^m(\sigma, \sigma^U) = A^T$ and $\mathcal{C}^{m-1}(\sigma, \sigma^U) = \bigoplus_{s \notin U} A^{T \cup \{s\}}$, so

$$\mathcal{H}^m(\sigma, \sigma^U) = A^T / \sum_{s \notin U} A^{T \cup \{s\}},$$

proving (7.1). (In particular, $\mathcal{H}^m(\sigma) = D^T$ and $\mathcal{H}^n(\Delta) = D^\emptyset$.) In the exact sequence (7.4), we have $\mathcal{H}^m(\sigma, \partial\sigma) = A^T$ and, by (7.1), $\mathcal{H}^m(\sigma, \sigma^{S-U}) = A^T / \sum_{s \in U} A^{T \cup \{s\}}$; hence, (7.2). Using (7.1) to calculate the second and third terms of (7.5), we get

$$\mathcal{H}^{m-1}(\sigma^U) = \sum_{s \in S-T} A^{T \cup \{s\}} / \sum_{s \in (S-T)-U} A^{T \cup \{s\}}$$

and this can be rewritten as (7.3). \square

By Proposition 7.1, D^T is the free abelian group $\mathcal{H}^m(\Delta_T)$. So, for each $T \in \mathcal{S}$, we can choose a splitting $\iota_T : D^T \rightarrow A^T$ of the projection map $A^T \rightarrow D^T$.

Definition 7.2. Let $\hat{A}^T := \iota_T(D^T)$. (It is a direct summand of the free abelian group A^T .)

Proposition 7.3.

$$\mathcal{H}^m(\sigma, \sigma^U) = \bigoplus_{\substack{V \supseteq T \\ V - T \subseteq U}} \hat{A}^V \quad (7.6)$$

$$\mathcal{H}^{m-1}(\sigma^U, \partial(\sigma^U)) = \bigoplus_{\substack{V \supseteq T \\ (V - T) \cap U \neq \emptyset}} \hat{A}^V \quad (7.7)$$

$$\mathcal{H}^{m-1}(\sigma^U) = \bigoplus_{\substack{V \supseteq T \\ V - T \subseteq U}} \hat{A}^V \quad (7.8)$$

Proof. Assume by induction that (7.6) through (7.8) hold for $\dim \sigma = m - 1$ and assume as well that they hold when $\dim \sigma = m$ and U is replaced by U' with $|U'| < |U|$. Write $U = \{s\} \cup U'$, for some $s \in U$. Consider the exact sequence of the triple $(\sigma^U, \sigma_s \cup \partial(\sigma^U), \partial(\sigma^U))$:

$$0 \rightarrow \mathcal{H}^{m-1}(\sigma^{U'}, \partial(\sigma^{U'})) \rightarrow \mathcal{H}^{m-1}(\sigma^U, \partial(\sigma^U)) \rightarrow \mathcal{H}^{m-1}(\sigma_s, (\sigma_s)^{S-U'}) \rightarrow 0, \quad (7.9)$$

where we have used the excisions $\mathcal{H}^*(\sigma^U, \sigma_s \cup \partial(\sigma^U)) = \mathcal{H}^*(\sigma^{U'}, \partial(\sigma^{U'}))$ and $\mathcal{H}^*(\sigma_s \cup \partial(\sigma^U), \partial(\sigma^U)) = \mathcal{H}^*(\sigma_s, (\sigma_s)^{S-U'})$ to rewrite the first and third terms. By induction,

$$\begin{aligned} \mathcal{H}^{m-1}(\sigma^{U'}, \partial(\sigma^{U'})) &= \bigoplus_{\substack{V \supseteq T \\ V \cap U' \neq \emptyset}} \hat{A}^V, \\ \mathcal{H}^{m-1}(\sigma_s, (\sigma_s)^{S-U'}) &= \bigoplus_{\substack{V \supseteq T \cup \{s\} \\ V \subseteq T \cup \{s\} \cup U'}} \hat{A}^V \end{aligned}$$

Substituting these into the last two terms of (7.9), we get

$$\mathcal{H}^{m-1}(\sigma^U, \partial(\sigma^U)) = \bigoplus_{\substack{V \supseteq T \\ V \cap U \neq \emptyset}} \hat{A}^V,$$

which is (7.7). Next consider the Mayer-Vietoris sequence of $\sigma^U = \sigma^{U'} \cup \sigma_s$:

$$0 \rightarrow \mathcal{H}^{m-2}((\sigma_s)^{U'}) \rightarrow \mathcal{H}^{m-1}(\sigma^U) \rightarrow \mathcal{H}^{m-1}(\sigma^{U'}) \oplus \mathcal{H}^{m-1}(\sigma_s) \rightarrow 0. \quad (7.10)$$

By induction,

$$\mathcal{H}^{m-2}((\sigma_s)^{U'}) = \bigoplus_{\substack{V \supseteq T \cup \{s\} \\ V - (T \cup \{s\}) \subseteq U'}} \hat{A}^V \quad \text{and} \quad \mathcal{H}^{m-1}(\sigma^{U'}) = \bigoplus_{\substack{V \supseteq T \\ V - T \subseteq U'}} \hat{A}^V$$

and $\mathcal{H}^{m-1}(\sigma_s) = \hat{A}^{T \cup \{s\}}$. Substituting these into (7.10) we get

$$\mathcal{H}^{m-1}(\sigma^U) = \bigoplus_{\substack{V \supseteq T \\ V - T \subseteq U}} \hat{A}^V,$$

which is (7.8). Sequence (7.5) is

$$0 \rightarrow \mathcal{H}^{m-1}(\sigma^U) \rightarrow \mathcal{H}^m(\sigma, \sigma^U) \rightarrow \mathcal{H}^m(\sigma) \rightarrow 0.$$

Substituting (7.8) for the first term and \hat{A}^T for the third, we get formula (7.6) for the middle term. \square

We have $\mathcal{H}^m(\sigma, \partial\sigma) = A^T$. Hence, in the special case $U = S - T$ formulas (7.6) and (7.8) give the following theorem (cf. [25], [11, Thm. 9.11], [10, Cor. 3.3]).

Theorem 7.4. (The Decomposition Theorem). *For each subset T of S*

$$A^T = \bigoplus_{V \supseteq T} \hat{A}^V.$$

For any \mathbf{Z} -submodule $B \subset A$, put $B^T := A^T \cap B$. Suppose we have a direct sum decomposition of \mathbf{Z} -modules, $A = B \oplus C$, so that for each $T \subseteq S$, $A^T = B^T \oplus C^T$. As explained in [10, §2] this leads to a decomposition of coefficient systems $\mathcal{I}(A) = \mathcal{I}(B) \oplus \mathcal{I}(C)$ so that for any mirrored CW complex X we have a decomposition of cochain complexes $\mathcal{C}^*(X; \mathcal{I}(A)) = \mathcal{C}^*(X; \mathcal{I}(B)) \oplus \mathcal{C}^*(X; \mathcal{I}(C))$. Since

$$(\hat{A}^V)^T = \begin{cases} \hat{A}^V & \text{if } V \supseteq T, \\ 0 & \text{otherwise,} \end{cases}$$

the formula in the Decomposition Theorem satisfies $(A^0)^T = \bigoplus_{V \supseteq T} (\hat{A}^V)^T$, for all $T \subseteq S$. So, we get a decomposition of coefficient systems $\mathcal{I}(A) = \bigoplus \hat{A}^V$ and a corresponding decomposition of cochain complexes:

$$\mathcal{C}^*(X; \mathcal{I}(A)) = \bigoplus_V \mathcal{C}^*(X; \mathcal{I}(\hat{A}^V)) \quad (7.11)$$

8 Cohomology of buildings

Just as in [10, Thm. 3.5], the Decomposition Theorem (Theorem 7.4) implies the following.

Theorem 8.1. (cf. [15, Cor. 8.2], [11, Thm. 10.3], [10, Thm. 3.5]). *Suppose X is a finite, mirrored CW complex. Then*

$$H_c^*(\mathcal{U}(\Phi, X^f)) \cong \bigoplus_{T \in \mathcal{S}} H^*(X, X^{S-T}) \otimes \hat{A}^T.$$

Proof. By Lemma 6.3, $H_c^*(\mathcal{U}(\Phi, X^f))$ is the cohomology of $\mathcal{C}^*(X; \mathcal{I}(A))$. Formula (7.11) gives a decomposition of cochain complexes:

$$\mathcal{C}^*(X; \mathcal{I}(A)) = \bigoplus_{T \in \mathcal{S}} \mathcal{C}^*(X; \mathcal{I}(\hat{A}^T))$$

We have

$$\mathcal{C}^k(X; \mathcal{I}(\hat{A}^T)) = \prod_{c \in X^{(k)}} \hat{A}^T \cap A^{S(c)} = \prod_{\substack{c \in X^{(k)} \\ c \notin X^{S-T}}} \hat{A}^T.$$

So, an element of $\mathcal{C}^k(X; \mathcal{I}(\hat{A}^T))$ is just an ordinary \hat{A}^T -valued cochain on X which vanishes on X^{S-T} , i.e.,

$$\mathcal{C}^*(X; \mathcal{I}(\hat{A}^T)) = C^*(X, X^{S-T}) \otimes \hat{A}^T; \quad \text{so,}$$

$$\mathcal{C}^*(X; \mathcal{I}(A)) = \bigoplus_{T \in \mathcal{S}} C^*(X, X^{S-T}) \otimes \hat{A}^T.$$

Taking cohomology, we get the result. \square

The most important special case of the previous theorem is the following.

Corollary 8.2.

$$H_c^*(\mathcal{U}(\Phi, K)) \cong \bigoplus_{T \in \mathcal{S}} H^*(K, K^{S-T}) \otimes \hat{A}^T.$$

9 The G -module structure on cohomology

We assume X has a W -finite mirror structure (i.e., $X = X^f$). Suppose G is a group of automorphisms of Φ . Then A , A^T , $A^{>T}$ and D^T are naturally right G -modules and so are the cochain complex $\mathcal{C}_c^*(\mathcal{U}(\Phi, X))$ and its cohomology groups. The discussion in Section 8 is well adapted to studying

the G -module structure of $H_c^*(\mathcal{U}(\Phi, X))$. As in [10], we should not expect a direct sum splitting of G -modules analogous nonequivariant splitting of Theorem 8.1; rather there should be a filtration of cohomology by G -modules with associated graded terms similar to those in the direct sum. As we indicate below, this is the case.

For each nonnegative integer p define a G -submodule F_p of A by

$$F_p := \sum_{|T| \leq p} A^T.$$

This gives a decreasing filtration:

$$A = F_0 \supset \cdots \supset F_p \supset F_{p+1} \cdots . \quad (9.1)$$

As in [10], it follows from the Decomposition Theorem that the associated graded terms are

$$F_p/F_{p+1} = \bigoplus_{|T|=p} D^T.$$

As in Section 6, we get a coefficient system $\mathcal{I}(F_p)$, a cochain complex $\mathcal{C}^*(X; \mathcal{I}(F_p))$ and corresponding cohomology groups $\mathcal{H}^*(X; \mathcal{I}(F_p))$. The filtration (9.1) leads to a filtration of $\mathcal{H}^*(X; \mathcal{I}(A)) (= H_c^*(\mathcal{U}(\Phi, X)))$ by right G -modules,

$$\mathcal{H}^*(X; \mathcal{I}(A)) \supset \cdots \supset \mathcal{H}^*(X; \mathcal{I}(F_p)) \supset \mathcal{H}^*(X; \mathcal{I}(F_{p+1})) \cdots . \quad (9.2)$$

As in [10], the Decomposition Theorem implies that

$$0 \longrightarrow \mathcal{H}^*(X; \mathcal{I}(F_{p+1})) \longrightarrow \mathcal{H}^*(X; \mathcal{I}(F_p)) \longrightarrow H^* \left(\frac{\mathcal{C}^*(X; \mathcal{I}(F_p))}{\mathcal{C}^*(X; \mathcal{I}(F_{p+1}))} \right) \longrightarrow 0$$

is short exact. From this we immediately deduce the following.

Theorem 9.1. *Suppose G is a group of automorphisms of Φ . Then the filtration (9.2) of $H_c^*(\mathcal{U}(\Phi, X))$ by right G -modules has associated graded terms,*

$$\mathcal{H}^*(X; F_p)/\mathcal{H}^*(X; F_{p+1}) \cong \bigoplus_{T \in \mathcal{S}^{(p)}} H^*(X, X^{S-T}) \otimes D^T.$$

Cocompact G -action. In this final paragraph we assume that there are only finitely many G -orbits on Φ and that X is a finite complex. These hypotheses imply that the quotient space $\mathcal{U}(\Phi, X)/G$ is compact and since X is a finite complex, each of the cohomology groups $H^*(X)$ is finitely generated. As a corollary to Theorem 9.1, we have the following.

Corollary 9.2. *With the above hypotheses, $H_c^*(\mathcal{U}(\Phi, X))$ is a finitely generated G -module.*

Proof. The assumption that G has only finitely many orbits of chambers implies that each A^T is finitely generated G -module; hence, so is each D^T . \square

Example 9.3. Suppose G is chamber transitive, i.e., suppose Φ consists of a single G -orbit. Choose a base chamber φ_0 , let B be its stabilizer and for each $T \in \mathcal{S}$, let G_T be the stabilizer of the T -residue containing φ_0 . Then A is the G -module $\mathbf{Z}(G/B)$ of finitely supported functions on G/B and A^T can be identified with $\mathbf{Z}(G/G_T)$. For each $U \supset T$, we have a natural inclusion $\mathbf{Z}(G/G_U) \rightarrow \mathbf{Z}(G/G_T)$ induced by the projection $G/G_T \rightarrow G/G_U$ and D^T is the G -module formed by dividing out the sum of the images of $\mathbf{Z}(G/G_U)$ in $\mathbf{Z}(G/G_T)$ for all $U \supset T$.

Here are some further corollaries to Corollary 8.2 and Theorem 9.1.

Corollary 9.4. *Suppose $\Gamma \subseteq G$ is a cocompact lattice in G . Then $H^*(\Gamma; \mathbf{Z}\Gamma)$ has, in filtration degree p , associated graded Γ -module*

$$\bigoplus_{T \in \mathcal{S}^{(p)}} H^*(K, K^{S-T}) \otimes D^T.$$

In particular, when Γ is torsion-free, its cohomological dimension is given by

$$\text{cd}(\Gamma) = \max\{k \mid H^k(K, K^{S-T}) \neq 0, \text{ for some } T \in \mathcal{S}\}.$$

A torsion-free group Γ is an n -dimensional duality group if $H^*(\Gamma; \mathbf{Z}\Gamma)$ is free abelian and concentrated in dimension n . Following [12, Definition 6.1], we say that the nerve of a Coxeter system has *punctured homology concentrated in dimension n* if for all $T \in \mathcal{S}$, $\tilde{H}^*(K^{S-T})$ is free abelian and concentrated in dimension n . Corollary 8.2 gives us a (correct) proof of the following result, stated in [12, Theorem 6.3].

Corollary 9.5. *Suppose $\Gamma \subseteq G$ is a torsion-free, cocompact lattice in G . Then the following are equivalent.*

- 1) Γ is an n -dimensional duality group.
- 2) W is an n -dimensional virtual duality group.
- 3) The nerve of (W, S) has punctured homology concentrated in dimension $n - 1$.

References

- [1] P. Abramenko and K. Brown, *Approaches to Buildings*, Springer, New York, 2008.
- [2] A. Borel and J.-P. Serre, *Cohomologie d'immeubles et de groupes S -arithmétiques*, *Topology* **15** (1976), 211–232.
- [3] N. Bourbaki, *Lie Groups and Lie Algebras, Chapters 4–6*, Springer, New York, 2002.
- [4] N. Brady, J. McCammond and J. Meier, *Local-to-asymptotic topology for cocompact CAT(0) complexes*, *Topology and its Applications* **131** (2003), 177–188.
- [5] M. Bridson and A. Haefliger, *Metric Spaces of Non-positive Curvature*, Springer, New York, 1999.
- [6] P.-E. Caprace and B. Rémy *Simplicité abstraite des groupes de Kac-Moody non affines*, *C. R. Acad. Sc. Paris* **342** (2006), 539–544.
- [7] M.W. Davis, *The cohomology of a Coxeter group with group ring coefficients*, *Duke Math. J.* **91** (1998), 297–314.
- [8] ———, *Buildings are CAT(0)*. In *Geometry and Cohomology in Group Theory (Durham, 1994)*, London Math. Soc. Lecture Note Ser. **252**, Cambridge Univ. Press, Cambridge, 1998, pp. 108–123.
- [9] ———, *The Geometry and Topology of Coxeter Groups*, London Math. Soc. Monograph Series, vol. 32, Princeton Univ. Press, 2007.
- [10] M.W. Davis, J. Dymara, T. Januszkiewicz and B. Okun, *Cohomology of Coxeter groups with group ring coefficients: II*, *Algebr. Geom. Topol.* **6** (2006), 1289–1318.
- [11] ———, *Weighted L^2 -cohomology of Coxeter groups*, *Geometry & Topology* **11** (2007), 47–138.
- [12] M.W. Davis and J. Meier, *The topology at infinity of Coxeter groups and buildings*, *Comment. Math. Helv.* **77** (2002), 746–766.
- [13] ———, *Erratum to “The topology at infinity of Coxeter groups and buildings”*, *Comment. Math. Helv.* **82** (2007), 235–236.
- [14] J. Dymara *Thin buildings*, *Geometry & Topology* **10** (2006), 667–694.

- [15] J. Dymara and T. Januszkiewicz, *Cohomology of buildings and their automorphism groups*, Invent. Math. **150** (2002), 579–627.
- [16] J. Dymara and D. Osajda, *Boundaries of right-angled hyperbolic buildings*, Fund. Math. **197** (2007), 123–165.
- [17] R. Geoghegan, *Topological Methods in Group Theory*, Springer, New York, 2008.
- [18] M. Gromov, *Hyperbolic groups* in *Essays in Group Theory*, edited by S. M. Gersten, M.S.R.I. Publ. 8, Springer, New York, 1987, pp. 75–264.
- [19] T. Januszkiewicz and J. Świątkowski, *Commensurability of graph products*, Algebr. Geom. Topol. **1** (2001), 587–603.
- [20] G. Moussong, *Hyperbolic Coxeter groups*, Ph.D. thesis, Ohio State University, 1988.
- [21] B. Rémy, *Topological simplicity, commensurator super-rigidity and nonlinearities of Kac-Moody groups. With an appendix by P. Bonvin*, Geom. Funct. Anal. **14** (2004), 810–852.
- [22] B. Rémy and M. Ronan, *Topological groups of Kac-Moody type, right-angled twinings and their lattices*, Comment. Math. Helv. **81** (2006), 191–219.
- [23] M. Ronan, *Lectures on Buildings*, Perspectives in Mathematics, vol. 7, Academic Press, San Diego, 1989.
- [24] B. Schulz, *Sphärische Unterkomplexe sphärischer Gebäude*, PhD thesis, Johann Wolfgang Goethe-Univ., Frankfurt a.M., 2005.
- [25] L. Solomon, *A decomposition of the group algebra of a finite Coxeter group*, J. of Algebra **9** (1968), 220–239.
- [26] A. Thomas, *Lattices acting on right-angled buildings*, Algebr. Geom. Topol. **6** (2006), 1215–1218.